19ème Journée d'automne

Suivi des maladies digestives et hépatobiliaires : collaboration médecins généralistes et spécialistes - qui fait quoi ?

Tests hépatiques perturbés

Ente Ospedaliero Cantonale

Andrea De Gottardi

Service de Gastroentérologie et Hépatologie, Ente Ospedaliero Cantonale,
Faculté de Médecine, Università della Svizzera Italiana, Lugano

Tests hépatiques

SYSTEM/FUNCTION	MARKER	LOCALISATION
Integrity of hepatocytes	AST	Liver, Muscle, Heart , Kidney, Pancreas, Lung
	ALT	Liver , Muscle, Heart, Kidney, Pancreas, Lung
Cholestasis	Alcaline phosphatase	Liver, Bone, Placenta
	GGT	With AP: Cholestasis Without AP: non hepatic origin

modif. Giannini, CMAJ, 2005

Transaminases, survie et mortalité

Transaminases élevées

ALT/AST level for increased mortality	Comments
AST>18	3X increase in all cause mortality
ALT 30-39	RR of liver mortality 2.9 (2.4-3.5) and 9.5 (7.9-11.5) in men, 3.8 (1.9-7.7) and 6.6 (1.5-25.6) in women
ALT 45-90 M 29-58 for F	SMR risk 1.32 for 1-2X ULN, and 1.78 for >2X ULN
ALT>30 for M ALT >19 for F	Increased liver related mortality

Kwo, Am J Gastro, 2017

Transaminases basses

Le Couteur, J Gerontol, 2010

Transaminases: cinétique

modif. Giannini, CMAJ, 2005

Cordoba, Hepatology 1998

Causes d'une perturbation chronique des tests hépatiques

Infection virale HBV, HCV, (HDV, HEV)	anti-HBc, anti-HBs, HBs-Ag anti-HCV, anti-HDV, HEV-RNA
Cholangite biliaire primitive	anti-mitochondries, IgM
Hépatite autoimmune	IgG, ANA, anti-muscle lisse, anti-LKM, anti-SLA
Hémochromatose, maladie de Wilson	ferritine, saturation de la transferrine, coeruloplasmine
Déficit en alpha-1 antitrypsine	alpha-1-AT
Médicaments (par ex. méthotrexate, statines)	???
MAFLD	???
Maladie alcoolique	CDT, éthylglucuronide

Phosphatase alcaline

Enzyme ubiquitaire, hautement exprimée dans le foie, dans les voies biliaires, le rein, l'os et le placenta.

Augmentation lors de cholestase suivant la GGT.

AUGMENTATION

Cholestase
 Hépatopathie aiguë ou chronique
 Maladies osseuses, Paget
 Hyperparathyroïdie
 Leucémie
 Lymphomes
 Sarcoïdose

DIMINUTION

Hypothyroïse
 Anémie
 Maladie de Wilson
 Contraceptifs oraux

Comment procéder?

Von Köckritz, Ther Umschau, 2017 & Modif. nach Giannini, CMAJ, 2005

Que faire quant tout est négatif?

- Revenir à l'histoire du patient: produits de phytothérapie, régime alimentaire, médicaments, hobbies, alcool...
- Refaire quelques tests: alpha1-antitrypsine, coeruloplasmine
- Considérer d'autres maladies: insuffisance cardiaque (droite), dysfonction thyroïdienne
- Chercher des maladies musculaires (CK, aldolase)
- Repeter l'imagerie
- Attendre....mais pas trop longtemps!

Que faire quant tout est négatif?

Fibrosis-4 (FIB-4) Index for Liver Fibrosis ☆

Noninvasive estimate of liver scarring in HCV and HBV patients, to assess need for biopsy.

When to Use ▼ Pearls/Pitfalls ▼ Why Use ▼

Age	49	years
AST Aspartate aminotransferase	34	U/L
Platelet count	275	$\times 10^9/L$ ↔
ALT Alanine aminotransferase	22	U/L

1.29 points

Fibrosis Stage: 0-1

FIB-4 scores <1.45 are 81% sensitive (NPV 90%) to rule out advanced fibrosis.

Copy Results 📄

Next Steps »»»

Investigations chez le spécialiste

Fibroscan

Shear wave elastography

Revenons à notre patiente

Ferritine sg	☐ (15 - 150) µg/l	1119 Δ H
Transferrine sg	☐ (24.7 - 44.4) µmol/l	17 Δ L
Coef. Sat. Transf. sg	☐ (0.15 - 0.35)	0.84 Δ H
CK sg	☐ (25 - 140) U/l	
LDH sg	☐ (135 - 214) U/l	388 Δ H
ASAT (GOT) sg	☐ (9 - 32) U/l	137 Δ H
ALAT (GPT) sg	☐ (9 - 36) U/l	34 Δ
Phosphatase alcaline sg	☐ (36 - 108) U/l	24 Δ L
Gamma-GT sg	☐ (6 - 42) U/l	191 Δ H
Bilirubine totale sg	☐ (0 - 21) µmol/l	99 Δ H
Bilirubine directe sg	☐ (0 - 10) µmol/l	55 Δ H

Low Serum Alkaline Phosphatase Activity in Wilson's Disease

WILLIAM A. SHAVER,† HARSHIKA BHATT AND BURTON COMBES

The Liver Unit, Department of Internal Medicine, The University of Texas Health Science Center at Dallas, Southwestern Medical School, Dallas, Texas 75235

Hepatology, 1986

Low serum alkaline phosphatase activity associated with severe Wilson's disease. Is the breakdown of alkaline phosphatase molecules caused by reactive oxygen species?

**Tadashi Hoshino*^a, Kazunari Kumasaka^a, Kinya Kawano^a,
Fujio Yamagishi^b, Iwao Koyama^c, Yohko Fujimori-Arai^c,
Takanori Nakajima^c, Tsugikazu Komoda*^c**

Fig. 2. Relative molecular size of the patient ALP by Western blotting using anti-bovine ALP serum. Lane 1, healthy child (ALP activity: 339 units/l); lane 2, healthy child (468 units/l); lane 3, the patient (9 units/l). kDa, kilodalton. The Western blotting conditions are described in Methods.

Clinica Chimica Acta, 1995

Rôle de la biopsie hépatique

H&E: noyaux avec glycogène

H&E: stéatose

Guindi, Semin Diagn Pathol, 2019

Rôle de la biopsie hépatique

Orcéine

Rhodanine

Guindi, Semin Diagn Pathol, 2019; Schilsky, Nat Rev Dis Prim, 2018

Schilsky, Nat Rev Dis Prim, 2018

Diagnostic

Test	Typical finding	False "negative"	False "positive"
Serum ceruloplasmin	Decreased by 50% of lower normal value	Normal levels in patients with marked hepatic inflammation Overestimation by immunologic assay Pregnancy, estrogen therapy	Low levels in: - malabsorption - aceruloplasminemia - heterozygotes
24-hour urinary copper	>1.6 $\mu\text{mol}/24\text{ h}$ >0.64 $\mu\text{mol}/24\text{ h}$ in children	Normal: - incorrect collection - children without liver disease	Increased: - hepatocellular necrosis - cholestasis - contamination
Serum "free" copper	>1.6 $\mu\text{mol}/\text{L}$	Normal if ceruloplasmin overestimated by immunologic assay	
Hepatic copper	>4 $\mu\text{mol}/\text{g}$ dry weight	Due to regional variation - in patients with active liver disease - in patients with regenerative nodules	Cholestatic syndromes
Kayser-Fleischer rings by slit lamp examination	Present	Absent - in up to 50% of patients with hepatic Wilson's disease - in most asymptomatic siblings	Primary biliary cirrhosis

EASL, CPG, 2012

Score de Leipzig

KF rings		Urinary copper in absence of acute hepatitis	
Present	2	Normal	0
Absent	0	1–2x ULN	1
Neurologic symptoms		> 2x ULN	2
Severe	2	Normal, but >5x ULN after D-penicillamine	2
Mild	1	Mutation Analysis	
Absent	0	On both chromosomes detected	4
Serum ceruloplasmin		On one chromosome detected	1
Normal (>200 mg/L)	0	No mutations detected	
100–200 mg/L	1		
<100 mg/L	2		
Coombs negative hemolytic anemia		KF, Kayser-Fleischer rings; ULN, upper limit of normal.	
Present	0	* if no quantitative liver copper available.	
Absent	1	Total score:	
Other tests		4 or more = diagnosis established	
Liver copper (in absence of cholestasis) (values provided as $\mu\text{mol}/\text{g}$; equivalent in $\mu\text{g}/\text{g}$ inserted)		3 = Diagnosis possible, more tests needed	
>5x upper limit of normal (>4 $\mu\text{mol}/\text{g}$ or >250 $\mu\text{g}/\text{g}$)	2	2 or less = Diagnosis very unlikely.	
0.8–4 $\mu\text{mol}/\text{g}$ (50–250 $\mu\text{g}/\text{g}$)	1		
Normal <0.8 $\mu\text{mol}/\text{g}$ (<50 $\mu\text{g}/\text{g}$)	-1		
Rhodanine positive granules	1*		

Ferenci, Liver Int, 2003

Principes du traitement

Drug	Therapeutic role	When developed	PRO	CON
<i>D-Penicillamine</i>	<i>Standard first-line treatment</i>	1956	<i>Very effective</i>	<ul style="list-style-type: none"> • <i>High rate of adverse effects (~ 30%, some severe)</i> • <i>Risk of neurological worsening*</i>
<i>Trientine (dihydrochloride)</i>	<i>Second-line treatment (BID or TID dosing); Candidate first-line treatment</i>	1969	<i>Very effective</i>	<ul style="list-style-type: none"> • <i>Some risk of neurological worsening</i> • <i>Cost</i>
<i>Zinc salts</i>	<i>Maintenance treatment; first-line treatment in selected clinical situations</i>	1980s	<ul style="list-style-type: none"> • <i>Effective</i> • <i>Little risk of neurological worsening</i> • <i>Additional metabolic benefits in hepatocytes?</i> 	<ul style="list-style-type: none"> • <i>Adherence difficult (TID, away from meals) → loss of disease control</i> • <i>Hepatic compromise indicated by elevated serum aminotransferases</i>

Roberts, Curr Gastroenterol Rep, 2018

Principes du traitement

D-pénicillamine (Metalcaptase, D)
 250 mg/j – 1500 mg/j
 Suppression médullaire
 Syndrome néphrotique
 Effets cutanés
 + pyridoxine 25-50 mg/sem

Trientine (Triogen, CH)
 250 mg/j – 1500 mg/j
 Effets cutanés
 Nausées
 Anémie

Zinc (Burgerstein gluconate de zinc, CH)
 30 mg/j
 Effets digestifs
 Augmentation amylase/lipase

Pendant la grossesse:
continuer le traitement

Allaitement:
Ne pas allaiter et ne pas interrompre le tt pour allaiter!

Rosencrantz, Semin Liv Dis, 2011

Take home messages: tests hépatiques perturbés

Le médecin de famille peut faire beaucoup
(s'il veut!)

Durée prolongée = consultation avec un spécialiste

Phosphatase alcaline normale ou diminuée:
pensez à la maladie de Wilson