

References used to establish recommendations for general palliative care practice

1. Australian Government Department of Health and Ageing. Guidelines for a Palliative Approach for Aged Care in the Community Setting — Best practice guidelines for the Australian context. Canberra: Australian Government Department of Health and Ageing. 2011.
2. Bernacki RE, Block SD, American College of Physicians High Value Care Task F. Communication about serious illness care goals: a review and synthesis of best practices. JAMA Internal Medicine. 2014;174(12):1994-2003.
3. Casey E, Hamilton H, Weir H. Guidelines for Palliative and End of Life Care in Nursing Homes and Residential Care Homes. Ireland: Guidelines and Audit implementation network. 2013.
4. Dans M, Smith T, Back A, Baker JN, Bauman JR, Beck AC, et al. NCCN Guidelines Insights: Palliative Care, Version 2.2017. Journal of the National Comprehensive Cancer Network : JNCCN. 2017;15(8):989-97.
5. Ferrell BR, Temel JS, Temin S, Alesi ER, Balboni TA, Basch EM, et al. Integration of Palliative Care Into Standard Oncology Care: American Society of Clinical Oncology Clinical Practice Guideline Update. J Clin Oncol. 2017;35(1):96-112.
6. Gaertner J, Siemens W, Meerpohl JJ, Antes G, Meffert C, Xander C, et al. Effect of specialist palliative care services on quality of life in adults with advanced incurable illness in hospital, hospice, or community settings: systematic review and meta-analysis. BMJ. 2017;357:j2925.
7. Gonella S, Basso I, Dimonte V, Martin B, Berchialla P, Campagna S, et al. Association Between End-of-Life Conversations in Nursing Homes and End-of-Life Care Outcomes: A Systematic Review and Meta-analysis. J Am Med Dir Assoc. 2019;20(3):249-61.
8. Guidelines and Protocols Advisory Committee. Palliative care for the patient with incurable cancer or advanced disease. Part 1: approach to care. Victoria: BCGuidelines.ca. 2017.
9. Guidelines and Protocols Advisory Committee. Palliative care for the patient with incurable cancer or advanced disease. Part 2: pain and symptom management. Victoria: Guidelines and Protocols Advisory Committee. 2017.
10. Haun MW, Estel S, Rucker G, Friederich HC, Villalobos M, Thomas M, et al. Early palliative care for adults with advanced cancer. Cochrane Database Systematic Review. 2017;6:CD011129.
11. Isenberg SR, Aslakson RA, Smith TJ. Implementing Evidence-Based Palliative Care Programs and Policy for Cancer Patients: Epidemiologic and Policy Implications of the 2016 American Society of Clinical Oncology Clinical Practice Guideline Update. Epidemiol Rev. 2017;39(1):123-31.
12. Federal Office of Public Health (FOPH), Swiss Conference of the Cantonal Ministers of Public Health (CMH), Palliative.ch. Framework Concept for Palliative Care in Switzerland. Bern. 2014.
13. Kernick LA, Hogg KJ, Millerick Y, Murtagh FEM, Djahit A, Johnson M. Does advance care planning in addition to usual care reduce hospitalisation for patients with advanced heart failure: A systematic review and narrative synthesis. Palliat Med. 2018;32(10):1539-51.
14. Low JTS, Rohde G, Pittordou K, Candy B, Davis S, Marshall A, et al. Supportive and palliative care in people with cirrhosis: International systematic review of the perspective of patients, family members and health professionals. J Hepatol. 2018;69(6):1260-73.
15. McCusker M, Ceronky L, Crone C, Epstein H, Greene B, Halvorson J, et al. Palliative care for adults. Bloomington: Insitute for Clinical Systems Improvement. 2013.
16. McLeod B. Hospice Palliative Care Symptom Assessment Guide and Guideline. In: Fraser South Health Region, editor. 5th ed: Fraser South Health Region. 2019.
17. Murray CD, McDonald C, Atkin H. The communication experiences of patients with palliative care needs: A systematic review and meta-synthesis of qualitative findings. Palliat Support Care. 2015;13(2):369-83.
18. National Coalition for Hospice and Palliative Care. Clinical practice guidelines for quality palliative care. Richmond: National Coalition for Hospice and Palliative Care. 2018
19. National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: Palliative Care. USA: National Comprehensive Cancer Network. 2019.
20. National Consensus Project for Quality Palliative Care. Clinical Practice Guidelines for Quality Palliative Care. 3rd ed. Pittsburgh: National Consensus Project for Quality Palliative Care. 2013.
21. Northern England Clinical Networks. Palliative and End of Life Care Guidelines. 4th ed. NHS. 2016.
22. Oczkowski SJ, Chung HO, Hanvey L, Mbuagbaw L, You JJ. Communication Tools for End-of-Life Decision-Making in Ambulatory Care Settings: A Systematic Review and Meta-Analysis. PLoS One. 2016;11(4):e0150671.
23. Palliative.ch. Soins palliatifs généraux: recommandations à l'attention des professionnels de santé et des prestataires de soins en médecine de premier recours en ambulatoire et en milieu stationnaire. Bern: Palliative.ch. 2019.
24. Palliative Care Australia. National Palliative Care Standards. 5th ed. Canberra: PCA. 2018.
25. Pringle J, Johnston B, Buchanan D. Dignity and patient-centred care for people with palliative care needs in the acute hospital setting: A systematic review. Palliat Med. 2015;29(8):675-94.
26. Rietjens JAC, Sudore RL, Connolly M, van Delden JJ, Drickamer MA, Droger M, et al. Definition and recommendations for advance care planning: an international consensus supported by the European Association for Palliative Care. Lancet Oncol. 2017;18(9):e543-e51.
27. The Gold Standards Framework Centre in End of Life Care CIC, Thomas K. The GSF pronostic indicator guidance, 4th edition. 2011.
28. Toye C, Tieman J, Oldham M, Abbey J, Currow D, Hegarty M, et al. Guidelines for a Palliative Approach for Aged Care in the Community Setting: Best practice guidelines for the Australian context. Curtin University of Technology, Curtin University. 2011.

ID-PALL[®] version 1.0, 2020

Instructions for use and glossary for the ID-PALL instrument
IDentification of patients requiring general or specialized PALLiative care.

&

Practice recommendations for general palliative care.

Document intended to be used by medical and
nursing teams non-specialized in palliative care.

Dr. Fabienne Teike Lüthi
PD Dr. Mathieu Bernard
PD Dr. Claudia Gamondi
Prof. Anne-Sylvie Ramelet
Prof. Gian Domenico Borasio



Introduction

ID-PALL is a tool that has been developed and validated to help you identify patients requiring general or specialized palliative care (PC). General PC is the care provided by non-specialized PC professionals, to patients with an incurable, progressive, life-limiting illness, but whose situation is stable, in order to help improve their quality of life. Specialized PC is provided by professionals specialized in palliative care, i.e., who have specific training and practice within a mobile team, a specialized consultation or a PC unit. Specialized care is intended for patients whose situation is unstable and/or complex, either due to difficulties assessing and managing physical symptoms, or due to acute levels of psychosocial and/or existential suffering. Specialized PC teams can also provide teams with help in developing specific assessment, communication and advance care planning skills, as well as support the writing of advance care planning documents with patients, or provide support when health care professionals feel helpless and experience difficulty.

It is therefore important to be able to quickly identify patients who need PC, and to be able to differentiate between those who need general PC and those who need specialized PC. In order to do this, the use of a screening instrument by health professionals has been recognized as beneficial.

Once a patient has been identified, it is also necessary for teams to have specific good practice recommendations to guide them in their daily practice.

Purpose


The purpose of this document is to :

- introduce ID-PALL and explain how to use it
- provide a glossary clarifying important terminology
- provide good practice recommendations for the clinical management of patients requiring general palliative care.

¹ Teike Lüthi F, Bernard M, Beauverd M, Gamondi C, Ramelet AS, Borasio GD. Identification of patients in need of general and specialised palliative care (ID-PALL®): item generation, content, and face validity of a new interprofessional screening instrument. BMC Palliat Care. 2020;19:1-11.

² Teike Lüthi F, Bernard M, Vanderlinden K, Ballabeni P, Gamondi C, Ramelet AS, Borasio GD. Measurement properties of ID-PALL, a new instrument for the IDentification of patients in need of general or specialised PALLiative care. JPSM. 2021.

³ La terminologie « patient » est utilisée à titre épïcène et regroupe également d'autres appellations telles que : résident, client, bénéficiaire de soins ou autre



ID-PALL® S : lexique

1. Presence of at least one severe and persistent symptom , including pain, that has not responded satisfactorily to treatment within 48 hours	Patients with palliative care needs commonly experience multiple symptoms. Pain is the most commonly identified symptom, but patients may experience dyspnea, nausea, fatigue, anxiety, or other sources of discomfort that do not respond to standard treatments. After 48 hours of unrelieved pain, it becomes clear that other strategies should be considered.
2. Difficulties in evaluating physical symptoms or psychological, social difficulties or spiritual distress	Difficulties may be encountered in assessing patients, either due to cognitive or linguistic difficulties, unfamiliarity with certain instruments, or difficulty in broacing certain topics. This can increase the level of complexity of the situation.
3. Disagreement or uncertainty on the part of the patient, people close to them or health professionals regarding, for example, medical treatments, resuscitation code or complex decisions	
4. The patient has severe psychosocial or existential suffering (eg: clinically relevant symptoms of anxiety or depression, feelings of isolation or being a burden, loss of sense of meaning or hope, desire to die, or has made a request for assisted suicide)	
5. People close to the patient experience severe psychosocial or existential suffering (eg: clinically relevant symptoms of anxiety or depression, major feelings of exhaustion, major disruption to the family system, loss of meaning or hope)	
6. Palliative sedation is envisaged (decreasing the level of consciousness using specific medication to relieve a refractory symptom)	A refractory symptom is a symptom that cannot be controlled to the patient's satisfaction despite properly conducted specialized palliative management. An intolerable symptom is determined by the patient as a symptom or condition that he or she can no longer tolerate.
7. Advance care planning or advance directives are difficult to establish with the patient or people close to them	
8. In your opinion, the patient, people close to them or health professionals could benefit from the intervention of palliative care specialists .	If it seems that you, the patient or their loved ones could use the support of palliative care specialists, but you feel that may not be able to express the need more explicitly, follow your intuition. Whether it's helping you cope with feelings of helplessness, arranging a complicated home-coming, supporting you through a difficult conversation, or assisting you with some care, palliative care specialists can be a resource.

Introduction and use of ID-PALL

Who assesses?

ID-PALL is a screening instrument that can be used by nurses and physicians.

When is it used?

ID-PALL should be completed within 48 hours of the patient’s admission or if any significant changes in their condition occur.

How does it work?

ID-PALL is divided into two parts. ID-PALL G is be filled out to identify general PS needs. If ID-PALL G is positive, ID-PALL S needs to be completed to identify specialized PC needs.

ID-PALL is scored as follows:

- For ID-PALL G, if the answer is ‘no’ to question 1 or ‘yes’ to at least one of statements 2, 3, or 4, the patient most likely requires general palliative care. In this case, please complete
- ID-PALL S next and refer to the recommendations for general palliative care practice.
- For ID-PALL S, if the answer is ‘yes’ to at least one of the statements, the patient would most likely benefit from a consultation with a specialized palliative care team.

ID-PALL is a screening tool. The purpose of this tool is to elicit discussions and build a therapeutic palliative care plan with the patient and an interprofessional team, based on the recommendations for general PC practice and, if necessary, with the support of a specialized PC team.

Below you will find:

- the ID-PALL instrument
- a glossary clarifying important concepts and terminology
- recommendations for general palliative care practice. These recommendations are to be implemented according to the patient and their family’s situation.

ID-PALL® G

Identification of patients in need of **General PALLiative** Care.
General palliative care is provided by professionals without specialised palliative care training in all care settings and contexts.
Please respond to all of the statements below relative to the patient's **current situation**:

Space reserved for patient's ID label

ID-PALL® S

Identification of patients in need of **Specialist PALLiative** Care.
Specialised palliative care is provided by or with professionals specialized in palliative care.
Please respond to all of the statements below, relative to the patient's **current situation, only when the response to the ID-PALL G is positive**:

1. Would you be surprised if this patient died in the next 12 months?	<input type="checkbox"/> Yes <input type="checkbox"/> No
2. The patient has a progressive illness or group of illnesses or comorbidities that limits their life expectancy AND presents (select all that are applicable): a decline in general functioning (with limited reversibility and an increase in need for support in day to day activities) OR a pronounced instability over the last 6 months (defined by: one uncontrolled symptom from the patient's point of view OR a pressure ulcer category ≥3 OR more than one acute delirium episode, infection, unscheduled hospitalisation or fall) OR psychosocial or existential suffering (of the patient or people close to them) OR the need for support in making decisions during the final stages of life	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Yes <input type="checkbox"/> No
3. Current or planned interruption of treatments with curative intent or vital support measures (eg: artificial ventilation, dialysis, artificial feeding or hydration)	<input type="checkbox"/> Yes <input type="checkbox"/> No
4. Request for comfort care or palliative care from the patient, people close to them or health professionals	<input type="checkbox"/> Yes <input type="checkbox"/> No

If you have ticked

NO to question 1 OR YES to at least ONE of the statements 2, 3, or 4,
the patient is likely to require general palliative care.
Please complete the ID-PALL S questionnaire on the next page and refer to the general palliative care practice recommendations.

1. Presence of at least one severe and persistent symptom , including pain, that has not responded satisfactorily to treatment within 48 hours.	<input type="checkbox"/> Yes <input type="checkbox"/> No
2. Difficulties in evaluating physical symptoms or psychological, social difficulties or spiritual distress	<input type="checkbox"/> Yes <input type="checkbox"/> No
3. Disagreement or uncertainty on the part of the patient, people close to them or health professionals regarding, for example, medical treatments, resuscitation code or complex decisions	<input type="checkbox"/> Yes <input type="checkbox"/> No
4. The patient has severe psychosocial or existential suffering (eg: marked symptoms of anxiety or depression, feelings of isolation or of being a burden, loss of meaning or hope, desire to die, or has made a request for assisted suicide)	<input type="checkbox"/> Yes <input type="checkbox"/> No
5. People close to the patient experience severe psychosocial or existential suffering (eg: marked symptoms of anxiety or depression, major feelings of exhaustion, major disruption to the functioning of the family system, loss of meaning or hope)	<input type="checkbox"/> Yes <input type="checkbox"/> No
6. Palliative sedation is envisaged (to relieve an intolerable refractory symptom by decreasing the level of consciousness using specific medication)	<input type="checkbox"/> Yes <input type="checkbox"/> No
7. Advance care plan or advance directives are difficult to establish with the patient and/or people close to them	<input type="checkbox"/> Yes <input type="checkbox"/> No
8. In your opinion, the patient, people close to them or health professionals could benefit from the intervention of palliative care specialists .	<input type="checkbox"/> Yes <input type="checkbox"/> No

If you have ticked

YES to ONE of the above statements, the patient is likely to require consultation of a specialist palliative care team.

ID-PALL® v1, 2020, F. Teike Lüthi et al.

ID-PALL® G : glossary

1. Would you be surprised if this patient died in the next 12 months ?

Described as «surprise question,» this question requires your intuitive response, based on what you know about the patient and their current situation.

2. The patient has a progressive illness or group of illnesses or comorbidities that **limits their life expectancy** **AND** presents:

This question is divided into two parts. When caring for a patient, the first question to ask oneself is whether the patient has a progressive life-limiting illness. Combining this criteria with one of the following elements will determine the need for general palliative care.

a decline in general functioning (with limited reversibility and an increased need for support in day to day activities)
(e.g., symptom of anxiety and depression; family, financial, cultural or life concerns; lack of meaning)
OR

Functional decline is a decrease in the ability to perform personal care, often associated with a deterioration in mobility and the ability to perform activities of daily living.

a pronounced instability over the last 6 months (defined by: one uncontrolled symptom from the patient's point of view OR a pressure ulcer category ≥3 OR more than one acute delirium episode, infection, unscheduled hospitalisation or fall)
OR
psychosocial or existential suffering (of the patient or people close to them)
OR

When a person's sense of completeness is compromised, he or she may feel unwell. This feeling may stem from multiple sources: psychological, social, spiritual and/or existential. In a holistic vision of the person, these dimensions are interrelated and influence each other.

the need for support in making decisions during the final stages of life

In some situations, patients and their loved ones may have difficulty making decisions about which treatments/ care to accept or where to go for care/end of life care, for example. Initiating Advance Care Planning can help them clarify their values and use these to guide them in their decision-making.

3. Current or planned interruption of treatments with curative intent or vital support measures (eg: artificial ventilation, dialysis, artificial feeding or hydration)

4. Request for comfort care or palliative care from the patient, people close to them or from health professionals